



Headteacher: Mrs H White

# Lubenham All Saints Primary School

School Lane, Lubenham

Market Harborough LE16 9TW

Tel: (01858) 462319 Fax: (01858) 433341

E-mail: [admin@lubenham.leics.sch.uk](mailto:admin@lubenham.leics.sch.uk)

22<sup>nd</sup> July 2017

Dear Parents/Carers

It has been brought to our attention that we have a case of slap cheek in school. For your information the symptoms and course of illness are detailed in the attached fact sheet.

Yours sincerely

Mrs Heather White  
Headteacher



## SLAP CHEEK SYNDROME

**Slapped cheek syndrome (also called fifth disease or parvovirus B19) is a viral infection that's most common in children, although it can affect people of any age. It usually causes a bright red rash on the cheeks.** Although the rash can look alarming, slapped cheek syndrome is normally a mild infection that clears up by itself in one to three weeks. Once you've had the infection, you're usually immune to it for life. However, slapped cheek syndrome can be more serious for some people. If you're pregnant, have a blood disorder or a weakened immune system and have been exposed to the virus, you should get medical advice.

### **Symptoms of slapped cheek syndrome**

Symptoms of slapped cheek syndrome usually develop 4-14 days after becoming infected, but sometimes may not appear for up to 21 days.

### **Initial symptoms**

Some people with slapped cheek syndrome won't notice any early symptoms, but most people will have the following symptoms for a few days:

- a slightly high temperature (fever) of around 38C (100.4F)
- a runny nose
- a [sore throat](#)
- a [headache](#)
- an upset stomach
- feeling generally unwell

The infection is most contagious during this initial period.

### **Slapped cheek rash**

After a few days, a distinctive bright red rash on both cheeks (the so-called "slapped cheeks") normally appears, although adults may not get this. By the time this rash develops, the condition is no longer contagious. After another few days, a light pink rash may also appear on the chest, stomach, arms and thighs. This often has a raised, lace-like appearance and may be itchy. The rashes will normally fade within a week or two, although occasionally the body rash may come and go for a few weeks after the infection has passed. This can be triggered by exercise, heat, anxiety or stress.

### **When to get medical advice**

You don't usually need to see your GP if you think you or your child has slapped cheek syndrome, as the condition normally gets better on its own.

## What to do if you or your child has slapped cheek

Slapped cheek syndrome is usually mild and should clear up without specific treatment.

If you or your child is feeling unwell, you can try the following to ease the symptoms:

- **rest and drink plenty of fluids** – babies should continue their normal feeds
- **for a fever, headaches or joint pain, you can take painkillers, such as [paracetamol](#) or [ibuprofen](#)** – children under 16 shouldn't take aspirin
- **to reduce itchiness, you can take [antihistamines](#) or use an [emollient \(moisturising lotion\)](#)** – some antihistamines are not suitable for young children, so check with your pharmacist first

Unless you or your child is feeling unwell, there's no need to stay away from school or work once the rash has developed, as the infection is no longer contagious by this point. It's a good idea to notify your child's school about the infection, so children who develop early symptoms can be spotted quickly and vulnerable people can be made aware that they may need to get medical advice.

## How do you get slapped cheek syndrome?

Slapped cheek syndrome is caused by a virus called parvovirus B19. This is found in the droplets in the coughs and sneezes of someone with the infection.

The virus is spread in a similar way to [colds](#) and [flu](#). You can become infected by:

- **inhaling droplets** that are sneezed or coughed out by someone with the infection
- **touching a contaminated surface or object** and then touching your mouth or nose

Someone with slapped cheek syndrome is infectious during the period before the rash develops. Once the rash appears, the condition can no longer be passed on.

## Preventing slapped cheek syndrome

It's very difficult to prevent slapped cheek syndrome, because people who have the infection are most contagious before they develop any obvious symptoms. However, making sure that everyone in your household washes their hands frequently may help stop the infection from spreading.

There's currently no vaccine available to protect you against the condition.